

- 1 a  $= \log_{10} \frac{3}{2}$   
 $= \log_{10} 3 - \log_{10} 2$   
 $= b - a$   
 b  $= \log_{10} (2^3 \times 3)$   
 $= 3 \log_{10} 2 + \log_{10} 3$   
 $= 3a + b$   
 c  $= \log_{10} (1.5 \times 100)$   
 $= \log_{10} 1.5 + \log_{10} 100$   
 $= b - a + 2$
- 2 a  $\log_3 x = \frac{5}{4}$   
 $x = 3^{\frac{5}{4}} = 3.95$  (3sf)  
 b  $3 \log_3 x - 5 \log_3 x = 4$   
 $\log_3 x = -2$   
 $x = 3^{-2} = \frac{1}{9}$
- 3 a i  $= \log_2 q^{\frac{1}{2}} = \frac{1}{2} \log_2 q = \frac{1}{2} p$   
 ii  $= \log_2 8 + \log_2 q = 3 + p$   
 b  $3 + p - \frac{1}{2} p = 2$   
 $p = \log_2 q = -2$   
 $\therefore q = 2^{-2} = \frac{1}{4}$
- 4  $2000 = 1000 \times 1.022^{4t}$   
 $2 = 1.022^{4t}$   
 $4t \lg 1.022 = \lg 2$   
 $t = \frac{\lg 2}{4 \lg 1.022} = 7.96$   
 $\therefore 8$  years
- 5 a  $(0, -3)$   
 b  $k = -4$   
 c  $(\frac{1}{3})^x - 4 = 0$   
 $(\frac{1}{3})^x = 4$   
 $x = \frac{\lg 4}{\lg \frac{1}{3}} = -1.26$  (3sf)
- 6 a  $\log_3 \frac{x+1}{x-2} = 1$   
 $\frac{x+1}{x-2} = 3$   
 $x+1 = 3x-6$   
 $x = \frac{7}{2}$   
 b  $(2x+1) \lg 3 = (x-4) \lg 2$   
 $x(\lg 2 - 2 \lg 3) = \lg 3 + 4 \lg 2$   
 $x = \frac{\lg 3 + 4 \lg 2}{\lg 2 - 2 \lg 3}$
- 7 a i  $= 2^{-1}(2^x) = \frac{1}{2} t$   
 ii  $= 2(2^{2x}) = 2(2^x)^2 = 2t^2$   
 b  $2t^2 - 7t + 6 = 0$   
 $(2t-3)(t-2) = 0$   
 $t = 2^x = \frac{3}{2}, 2$   
 $x = \frac{\lg \frac{3}{2}}{\lg 2}, 1 = 0.585$  (3sf), 1
- 8 a  $\log_2 (3x+5) + 3 = 7$   
 $3x+5 = 2^4 = 16$   
 $x = \frac{11}{3}$   
 b  $\log_2 (x+1) + \log_2 (3x-1) = 5$   
 $(x+1)(3x-1) = 2^5 = 32$   
 $3x^2 + 2x - 33 = 0$   
 $(3x+11)(x-3) = 0$   
 $x = -\frac{11}{3}, 3$   
 for real  $\log_2 (3x-1), x > \frac{1}{3} \therefore x = 3$

9 a  $x + 4 = \frac{5}{4}x$

$x = 16$

b  $y + 2 = \frac{12}{y+1}$

$(y + 2)(y + 1) = 12$

$y^2 + 3y - 10 = 0$

$(y + 5)(y - 2) = 0$

$y > 0 \therefore y = 2$

c  $\log_y x = \log_2 16 = 4$

10 a  $t = 0 \Rightarrow n = 2000$

b  $3600 = \frac{18000}{1+8c^{-3}}$

$1 + 8c^{-3} = 5$

$c^{-3} = \frac{1}{2}$

$c^3 = 2$

$c = \sqrt[3]{2}$

c  $4000 = \frac{18000}{1+8c^{-t}}$

$1 + 8c^{-t} = \frac{9}{2}$

$c^{-t} = \frac{7}{16}$

$-t = \frac{\lg \frac{7}{16}}{\lg \sqrt[3]{2}}$

$t = 3.578 \text{ weeks} = 25 \text{ days}$

11 a i  $\log_8 x^2 = 2 \log_8 x = 2y$

ii  $y = \log_8 x \Rightarrow x = 8^y = 2^{3y}$

$\therefore \log_2 x = 3y$

b  $3(2y) + 3y = 6$

$y = \log_8 x = \frac{2}{3}$

$\therefore x = 8^{\frac{2}{3}} = 4$

12  $\log_2 y - \log_2 (3 - 2x) = 1 \Rightarrow \frac{y}{3-2x} = 2$

$\Rightarrow y = 6 - 4x$

$\log_4 xy = \frac{1}{2} \Rightarrow xy = 4^{\frac{1}{2}} = 2$

sub.  $x(6 - 4x) = 2$


$2x^2 - 3x + 1 = 0$

$(2x - 1)(x - 1) = 0$

$x = \frac{1}{2}, 1$

$\therefore x = \frac{1}{2}, y = 4 \text{ or } x = 1, y = 2$

13 a



b at A,  $2^x + 1 = (\frac{1}{2})^x$

$(2^x)^2 + 2^x = 1$

$2^{2x} + 2^x - 1 = 0$

c  $2^x = \frac{-1 \pm \sqrt{1+4}}{2}$

$2^x = \frac{-1 - \sqrt{5}}{2}$  [no sols] or  $\frac{-1 + \sqrt{5}}{2}$

$\therefore 2^x = \frac{1}{2} \sqrt{5} - \frac{1}{2}$

$\therefore y = (\frac{1}{2} \sqrt{5} - \frac{1}{2}) + 1 = \frac{1}{2}(\sqrt{5} + 1)$

14 a when  $x = 1$ ,

LHS =  $8 - 4(4) + 2 + 6 = 0$

$\therefore x = 1$  is a solution

b  $2^{3x} = (2^x)^3 = u^3$

$2^{2x} = (2^x)^2 = u^2$

$\therefore$  (I)  $\Rightarrow u^3 - 4u^2 + u + 6 = 0$

c  $x = 1 \Rightarrow u = 2 \therefore (u - 2)$  is a factor

$$\begin{array}{r} u^2 - 2u - 3 \\ u - 2 \overline{) u^3 - 4u^2 + u + 6} \\ \underline{u^3 - 2u^2} \phantom{+ u + 6} \\ - 2u^2 + u \phantom{+ 6} \\ \underline{- 2u^2 + 4u} \phantom{+ 6} \\ - 3u + 6 \\ \underline{- 3u + 6} \\ 0 \end{array}$$

$(u - 2)(u^2 - 2u - 3) = 0$

$(u - 2)(u - 3)(u + 1) = 0$

$u = 2^x = -1$  [no sols], 2 or 3

$x = 1$  (given) or  $\frac{\lg 3}{\lg 2} = 1.58$