

- 1 Calculate
- a**  $(\mathbf{i} + 2\mathbf{j}) \cdot (3\mathbf{i} + \mathbf{j})$                       **b**  $(4\mathbf{i} - \mathbf{j}) \cdot (3\mathbf{i} + 5\mathbf{j})$                       **c**  $(\mathbf{i} - 2\mathbf{j}) \cdot (-5\mathbf{i} - 2\mathbf{j})$
- 2 Show that the vectors  $(\mathbf{i} + 4\mathbf{j})$  and  $(8\mathbf{i} - 2\mathbf{j})$  are perpendicular.
- 3 Find in each case the value of the constant  $c$  for which the vectors  $\mathbf{u}$  and  $\mathbf{v}$  are perpendicular.
- a**  $\mathbf{u} = \begin{pmatrix} 3 \\ -1 \end{pmatrix}$ ,  $\mathbf{v} = \begin{pmatrix} c \\ 3 \end{pmatrix}$                       **b**  $\mathbf{u} = \begin{pmatrix} 2 \\ 1 \end{pmatrix}$ ,  $\mathbf{v} = \begin{pmatrix} 3 \\ c \end{pmatrix}$                       **c**  $\mathbf{u} = \begin{pmatrix} 2 \\ -5 \end{pmatrix}$ ,  $\mathbf{v} = \begin{pmatrix} c \\ -4 \end{pmatrix}$
- 4 Find, in degrees to 1 decimal place, the angle between the vectors
- a**  $(4\mathbf{i} - 3\mathbf{j})$  and  $(8\mathbf{i} + 6\mathbf{j})$                       **b**  $(7\mathbf{i} + \mathbf{j})$  and  $(2\mathbf{i} + 6\mathbf{j})$                       **c**  $(4\mathbf{i} + 2\mathbf{j})$  and  $(-5\mathbf{i} + 2\mathbf{j})$
- 5 Relative to a fixed origin  $O$ , the points  $A$ ,  $B$  and  $C$  have position vectors  $(9\mathbf{i} + \mathbf{j})$ ,  $(3\mathbf{i} - \mathbf{j})$  and  $(5\mathbf{i} - 2\mathbf{j})$  respectively. Show that  $\angle ABC = 45^\circ$ .
- 6 Calculate
- a**  $(\mathbf{i} + 2\mathbf{j} + 4\mathbf{k}) \cdot (3\mathbf{i} + \mathbf{j} + 2\mathbf{k})$                       **b**  $(6\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}) \cdot (\mathbf{i} - 3\mathbf{j} - \mathbf{k})$   
**c**  $(-5\mathbf{i} + 2\mathbf{k}) \cdot (\mathbf{i} + 4\mathbf{j} - 3\mathbf{k})$                       **d**  $(3\mathbf{i} + 2\mathbf{j} - 8\mathbf{k}) \cdot (-\mathbf{i} + 11\mathbf{j} - 4\mathbf{k})$   
**e**  $(3\mathbf{i} - 7\mathbf{j} + \mathbf{k}) \cdot (9\mathbf{i} + 4\mathbf{j} - \mathbf{k})$                       **f**  $(7\mathbf{i} - 3\mathbf{j}) \cdot (-3\mathbf{j} + 6\mathbf{k})$
- 7 Given that  $\mathbf{p} = 2\mathbf{i} + \mathbf{j} - 3\mathbf{k}$ ,  $\mathbf{q} = \mathbf{i} + 5\mathbf{j} - \mathbf{k}$  and  $\mathbf{r} = 6\mathbf{i} - 2\mathbf{j} - 3\mathbf{k}$ ,
- a** find the value of  $\mathbf{p} \cdot \mathbf{q}$ ,  
**b** find the value of  $\mathbf{p} \cdot \mathbf{r}$ ,  
**c** verify that  $\mathbf{p} \cdot (\mathbf{q} + \mathbf{r}) = \mathbf{p} \cdot \mathbf{q} + \mathbf{p} \cdot \mathbf{r}$
- 8 Simplify
- a**  $\mathbf{p} \cdot (\mathbf{q} + \mathbf{r}) + \mathbf{p} \cdot (\mathbf{q} - \mathbf{r})$                       **b**  $\mathbf{p} \cdot (\mathbf{q} + \mathbf{r}) + \mathbf{q} \cdot (\mathbf{r} - \mathbf{p})$
- 9 Show that the vectors  $(5\mathbf{i} - 3\mathbf{j} + 2\mathbf{k})$  and  $(3\mathbf{i} + \mathbf{j} - 6\mathbf{k})$  are perpendicular.
- 10 Relative to a fixed origin  $O$ , the points  $A$ ,  $B$  and  $C$  have position vectors  $(3\mathbf{i} + 4\mathbf{j} - 6\mathbf{k})$ ,  $(\mathbf{i} + 5\mathbf{j} - 2\mathbf{k})$  and  $(8\mathbf{i} + 3\mathbf{j} + 2\mathbf{k})$  respectively. Show that  $\angle ABC = 90^\circ$ .
- 11 Find in each case the value or values of the constant  $c$  for which the vectors  $\mathbf{u}$  and  $\mathbf{v}$  are perpendicular.
- a**  $\mathbf{u} = (2\mathbf{i} + 3\mathbf{j} + \mathbf{k})$ ,  $\mathbf{v} = (c\mathbf{i} - 3\mathbf{j} + \mathbf{k})$                       **b**  $\mathbf{u} = (-5\mathbf{i} + 3\mathbf{j} + 2\mathbf{k})$ ,  $\mathbf{v} = (c\mathbf{i} - \mathbf{j} + 3c\mathbf{k})$   
**c**  $\mathbf{u} = (c\mathbf{i} - 2\mathbf{j} + 8\mathbf{k})$ ,  $\mathbf{v} = (c\mathbf{i} + c\mathbf{j} - 3\mathbf{k})$                       **d**  $\mathbf{u} = (3c\mathbf{i} + 2\mathbf{j} + c\mathbf{k})$ ,  $\mathbf{v} = (5\mathbf{i} - 4\mathbf{j} + 2c\mathbf{k})$
- 12 Find the exact value of the cosine of the angle between the vectors
- a**  $\begin{pmatrix} 1 \\ 2 \\ -2 \end{pmatrix}$  and  $\begin{pmatrix} 8 \\ 1 \\ -4 \end{pmatrix}$                       **b**  $\begin{pmatrix} 4 \\ 1 \\ -2 \end{pmatrix}$  and  $\begin{pmatrix} -2 \\ 3 \\ -6 \end{pmatrix}$                       **c**  $\begin{pmatrix} 1 \\ 2 \\ -1 \end{pmatrix}$  and  $\begin{pmatrix} 1 \\ -7 \\ 2 \end{pmatrix}$                       **d**  $\begin{pmatrix} 5 \\ -3 \\ 4 \end{pmatrix}$  and  $\begin{pmatrix} 3 \\ -4 \\ -1 \end{pmatrix}$
- 13 Find, in degrees to 1 decimal place, the angle between the vectors
- a**  $(3\mathbf{i} - 4\mathbf{k})$  and  $(7\mathbf{i} - 4\mathbf{j} + 4\mathbf{k})$                       **b**  $(2\mathbf{i} - 6\mathbf{j} + 3\mathbf{k})$  and  $(\mathbf{i} - 3\mathbf{j} - \mathbf{k})$   
**c**  $(6\mathbf{i} - 2\mathbf{j} - 9\mathbf{k})$  and  $(3\mathbf{i} + \mathbf{j} + 4\mathbf{k})$                       **d**  $(\mathbf{i} + 5\mathbf{j} - 3\mathbf{k})$  and  $(-3\mathbf{i} - 4\mathbf{j} + 2\mathbf{k})$

- 14 The points  $A(7, 2, -2)$ ,  $B(-1, 6, -3)$  and  $C(-3, 1, 2)$  are the vertices of a triangle.
- Find  $\overrightarrow{BA}$  and  $\overrightarrow{BC}$  in terms of  $\mathbf{i}$ ,  $\mathbf{j}$  and  $\mathbf{k}$ .
  - Show that  $\angle ABC = 82.2^\circ$  to 1 decimal place.
  - Find the area of triangle  $ABC$  to 3 significant figures.
- 15 Relative to a fixed origin, the points  $A$ ,  $B$  and  $C$  have position vectors  $(3\mathbf{i} - 2\mathbf{j} - \mathbf{k})$ ,  $(4\mathbf{i} + 3\mathbf{j} - 2\mathbf{k})$  and  $(2\mathbf{i} - \mathbf{j})$  respectively.
- Find the exact value of the cosine of angle  $BAC$ .
  - Hence show that the area of triangle  $ABC$  is  $3\sqrt{2}$ .
- 16 Find, in degrees to 1 decimal place, the acute angle between each pair of lines.
- $\mathbf{r} = \begin{pmatrix} 1 \\ 3 \\ -1 \end{pmatrix} + \lambda \begin{pmatrix} 4 \\ -4 \\ 2 \end{pmatrix}$  and  $\mathbf{r} = \begin{pmatrix} 5 \\ -2 \\ 1 \end{pmatrix} + \mu \begin{pmatrix} 8 \\ 0 \\ -6 \end{pmatrix}$
  - $\mathbf{r} = \begin{pmatrix} 0 \\ -3 \\ 7 \end{pmatrix} + \lambda \begin{pmatrix} 6 \\ -1 \\ -18 \end{pmatrix}$  and  $\mathbf{r} = \begin{pmatrix} 4 \\ 6 \\ -3 \end{pmatrix} + \mu \begin{pmatrix} 4 \\ -12 \\ 3 \end{pmatrix}$
  - $\mathbf{r} = \begin{pmatrix} 7 \\ 1 \\ 5 \end{pmatrix} + \lambda \begin{pmatrix} 1 \\ -1 \\ 3 \end{pmatrix}$  and  $\mathbf{r} = \begin{pmatrix} -2 \\ 6 \\ -3 \end{pmatrix} + \mu \begin{pmatrix} 2 \\ -5 \\ 3 \end{pmatrix}$
  - $\mathbf{r} = \begin{pmatrix} 2 \\ -3 \\ -9 \end{pmatrix} + \lambda \begin{pmatrix} -4 \\ -6 \\ 7 \end{pmatrix}$  and  $\mathbf{r} = \begin{pmatrix} 11 \\ 1 \\ -2 \end{pmatrix} + \mu \begin{pmatrix} 5 \\ -1 \\ -8 \end{pmatrix}$
- 17 Relative to a fixed origin, the points  $A$  and  $B$  have position vectors  $(5\mathbf{i} + 8\mathbf{j} - \mathbf{k})$  and  $(6\mathbf{i} + 5\mathbf{j} + \mathbf{k})$  respectively.
- Find a vector equation of the straight line  $l_1$  which passes through  $A$  and  $B$ .  
The line  $l_2$  has the equation  $\mathbf{r} = 4\mathbf{i} - 3\mathbf{j} + 5\mathbf{k} + \mu(-5\mathbf{i} + \mathbf{j} - 2\mathbf{k})$ .
  - Show that lines  $l_1$  and  $l_2$  intersect and find the position vector of their point of intersection.
  - Find, in degrees, the acute angle between lines  $l_1$  and  $l_2$ .
- 18 Find, in degrees to 1 decimal place, the acute angle between the lines with cartesian equations
- $$\frac{x-2}{3} = \frac{y}{2} = \frac{z+5}{-6} \quad \text{and} \quad \frac{x-4}{-4} = \frac{y+1}{7} = \frac{z-3}{-4}.$$
- 19 The line  $l$  has the equation  $\mathbf{r} = 7\mathbf{i} - 2\mathbf{k} + \lambda(2\mathbf{i} - \mathbf{j} + 2\mathbf{k})$  and the line  $m$  has the equation  $\mathbf{r} = -4\mathbf{i} + 7\mathbf{j} - 6\mathbf{k} + \mu(5\mathbf{i} - 4\mathbf{j} - 2\mathbf{k})$ .
- Find the coordinates of the point  $A$  where lines  $l$  and  $m$  intersect.
  - Find, in degrees, the acute angle between lines  $l$  and  $m$ .  
The point  $B$  has coordinates  $(5, 1, -4)$ .
  - Show that  $B$  lies on the line  $l$ .
  - Find the distance of  $B$  from  $m$ .
- 20 Relative to a fixed origin  $O$ , the points  $A$  and  $B$  have position vectors  $(9\mathbf{i} + 6\mathbf{j})$  and  $(11\mathbf{i} + 5\mathbf{j} + \mathbf{k})$  respectively.
- Show that for all values of  $\lambda$ , the point  $C$  with position vector  $(9 + 2\lambda)\mathbf{i} + (6 - \lambda)\mathbf{j} + \lambda\mathbf{k}$  lies on the straight line  $l$  which passes through  $A$  and  $B$ .
  - Find the value of  $\lambda$  for which  $OC$  is perpendicular to  $l$ .
  - Hence, find the position vector of the foot of the perpendicular from  $O$  to  $l$ .
- 21 Find the coordinates of the point on each line which is closest to the origin.
- $\mathbf{r} = -4\mathbf{i} + 2\mathbf{j} + 7\mathbf{k} + \lambda(\mathbf{i} + 3\mathbf{j} - 4\mathbf{k})$
  - $\mathbf{r} = 7\mathbf{i} + 11\mathbf{j} - 9\mathbf{k} + \lambda(6\mathbf{i} - 9\mathbf{j} + 3\mathbf{k})$